Delayed mode quality control of MOCCA Argo float 3901948

Kamila Walicka

British Oceanographic Data Centre (BODC), National Oceanography Centre Joseph Proudman Building, 6, Brownlow St, Liverpool L3 5DA

February 7, 2020

Summary

The cycles from 1 to 55 well reflect the Argo referenced data, the associated error 0.01, QC=1. The CTD data after cycle around 30 showed are very poor distributed. From cycle 56 to 77 there has been detected drift. The OWC correction has been applied, QC=1, error 0.014.

WMO number	DM correction
3901948	Drift detected

Table 1: Correction applied in delayed mode.

Contents

1	Introduction	3
2	Quality Check of Argo Float Data 2.1 Time Series of Vertical Distribution of Data 2.2 Comparison between Argo Float and Climatology 2.3 Satellite Altimeter comparison	3 3 4 8
3	Correction of Salinity Data 3.1 Comparison between Argo floats and CTD Climatlogy 3.1.1 Configuration 3.1.2 Results 3.2 Comparison between Argo floats and Argo Climatlogy 3.2.1 Configuration 3.2.2 Results	8 8 12 20 20 24
4	Final Checks	აა 33

1 Introduction

Delayed mode analysis was performed for float number 470894i (3901948) where salinity and temperature values were separately compared to nearby historical CTD profiles and nearby Argo profiles as a reference database. The OWC (Cabanes et al., 2016) method was run to estimate a salinity offset and/or a salinity drift. For more information about float 470894i (3901948) click on the following link: http://www.ifremer.fr/argoMonitoring/float/3901948

2 Quality Check of Argo Float Data

2.1 Time Series of Vertical Distribution of Data



Float 3901948 Potential Temperature

Figure 1: Float 3901948. Time series of the vertical distribution of potential temperature (°C).



Figure 2: Float 3901948. Time series of the vertical distribution of practical salinity (PSU).

2.2 Comparison between Argo Float and Climatology

The comparison between float 3901948 and data from WMO boxes $+/-10^{\circ}$ of latitude and longitude shows that the Argo profiles fit within the expected ranges (Figures 3, 4 and 5). This result confirms that float 3901948 represents relatively stable and consistent with the expected physical conditions in this region.



Figure 3: Float 3901948. Float profile of potential temperature (°C) plotted with climatology from the spatial range of 10 °. The black and blue cycles indicates the first and the last Argo profile, respectively. Green symbols represent other Argo profiles.



Figure 4: Float 3901948. Float profile of salinity (dimensionless) plotted with climatology from the spatial range of 10 °. The black and blue cycles indicates the first and the last Argo profile, respectively. Green symbols represent other Argo profiles.



Figure 5: Float 3901948. Theta/S plotted with climatology from the spatial range of 10 °. The black and blue cycles indicates the first and the last Argo profile, respectively. Green symbols represent other Argo profiles.

2.3 Satellite Altimeter comparison



Figure 6: Float 3901948. The comparison betweeen the Sea Surface Height(SSH) from the satellite altimetry and Dynamic Height Anomaly(DHA)extracted from the Argo float temperature and salinity data

3 Correction of Salinity Data

3.1 Comparison between Argo floats and CTD Climatlogy

3.1.1 Configuration

% -----%

```
%
    Climatology Data Input Paths
%
HISTORICAL_DIRECTORY=/users/argo/climatology
HISTORICAL_CTD_PREFIX=/historical_ctd/CTD_for_DMQC_2019V01/ctd_
HISTORICAL_BOTTLE_PREFIX=/historical_bot/WOD2001_v2/bot_
HISTORICAL_ARGO_PREFIX=/argo_profiles/ARGO_for_DMQC_2019V03/argo_
%
%
    Float Input Path
%
FLOAT_SOURCE_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_source/
FLOAT_SOURCE_POSTFIX=.mat
%
%
    Mapping Output Path
%
FLOAT_MAPPED_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_mapped/ctd/
FLOAT_MAPPED_PREFIX=map_
FLOAT_MAPPED_POSTFIX=.mat
%
%
    Calibration Output Path
%
FLOAT_CALIB_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_calib/ctd/
FLOAT_CALIB_PREFIX=cal_
FLOAT_CALSERIES_PREFIX=calseries_
FLOAT_CALIB_POSTFIX=.mat
%
%
    Diagnostic Plots Output Path
%
FLOAT_PLOTS_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_plots/ctd/
%
```

```
% Constants File Path
%
```

CONFIG_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/constants/ CONFIG_COASTLINES=coastdat.mat CONFIG_WMO_BOXES=wmo_boxes_ctd.mat CONFIG_SAF=TypicalProfileAroundSAF.mat % % max number of historical casts used in objective mapping CONFIG_MAX_CASTS=310 % 1=use PV constraint, 0=don't use PV constraint, in objective mapping MAP_USE_PV=1 % 1=use SAF separation criteria, 0=don't use SAF separation criteria, in objective mapping MAP_USE_SAF=1 % spatial decorrelation scales, in degrees MAPSCALE_LONGITUDE_LARGE=6 MAPSCALE_LONGITUDE_SMALL=3 MAPSCALE_LATITUDE_LARGE=4 MAPSCALE_LATITUDE_SMALL=2 % cross-isobath scales, dimensionless, see BS(2005) MAPSCALE_PHI_LARGE=0.1 MAPSCALE_PHI_SMALL=0.02 % temporal decorrelation scale, in years MAPSCALE_AGE=5 MAPSCALE_AGE_LARGE=10 % exclude the top xxx dbar of the water column MAP_P_EXCLUDE=100 % only use historical data that are within +/- yyy dbar from float data MAP_P_DELTA=200



Figure 7: Float 3901948. Trajectory of the Argo float plotted with historical CTD data (blue dots). The black contours indicate continent contours.



3901948 uncalibrated float data (-) and mapped salinity (o) with objective errors

Figure 8: Float 3901948. T/S diagram for the uncalibrated float data and mapped salinity. The collared circles indicate the theta levels with an objective errors. Diagram is showing every 3rd Argo profile.



3901948 potential conductivity (mmho/cm) multiplicative correction r with errors



Figure 9: Float 3901948. Time series of profiles of the potential conductivity (top) and vertically averaged salinity (bottom) with errors.



3901948 calibrated float data (-) and mapped salinity (o) with objective errors





Figure 11: Float 3901948. Salinity anomaly on θ levels.





Figure 12: Float 3901948. Salinities with errors on θ levels.



Figure 13: Float 3901948. Calibrated salinity anomaly on θ levels.



Figure 14: Float 3901948. Salinity, salinity variance on theta and OW chosen levels.

3.2 Comparison between Argo floats and Argo Climatlogy

3.2.1 Configuration

```
%
%
    Climatology Data Input Paths
%
HISTORICAL_DIRECTORY=/users/argo/climatology
HISTORICAL_CTD_PREFIX=/historical_ctd/CTD_for_DMQC_2019V01/ctd_
HISTORICAL_BOTTLE_PREFIX=/historical_bot/bot_
HISTORICAL_ARGO_PREFIX=/argo_profiles/ARGO_for_DMQC_2019V03/argo_
%
%
    Float Input Path
%
FLOAT_SOURCE_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_source/
FLOAT_SOURCE_POSTFIX=.mat
%
    Mapping Output Path
%
%
FLOAT_MAPPED_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_mapped/argo/
FLOAT_MAPPED_PREFIX=map_
FLOAT_MAPPED_POSTFIX=.mat
%
%
    Calibration Output Path
%
FLOAT_CALIB_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_calib/argo/
FLOAT_CALIB_PREFIX=cal_
FLOAT_CALSERIES_PREFIX=calseries_
FLOAT_CALIB_POSTFIX=.mat
%
%
    Diagnostic Plots Output Path
%
```

FLOAT_PLOTS_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_plots/argo/ % % Constants File Path % CONFIG_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/constants/ CONFIG_COASTLINES=coastdat.mat CONFIG_WMO_BOXES=wmo_boxes_argo.mat CONFIG_SAF=TypicalProfileAroundSAF.mat % % Objective Mapping Parameters % % max number of historical casts used in objective mapping CONFIG_MAX_CASTS=310 % 1=use PV constraint, 0=don't use PV constraint, in objective mapping MAP_USE_PV=1 % 1=use SAF separation criteria, 0=don't use SAF separation criteria, in objective mapping MAP_USE_SAF=1 % spatial decorrelation scales, in degrees MAPSCALE_LONGITUDE_LARGE=6 MAPSCALE_LONGITUDE_SMALL=3 MAPSCALE_LATITUDE_LARGE=4 MAPSCALE_LATITUDE_SMALL=2 % cross-isobath scales, dimensionless, see BS(2005) MAPSCALE_PHI_LARGE=0.1 MAPSCALE_PHI_SMALL=0.02 % temporal decorrelation scale, in years MAPSCALE_AGE=5 MAPSCALE_AGE_LARGE=10 % exclude the top xxx dbar of the water column MAP_P_EXCLUDE=100 % only use historical data that are within +/- yyy dbar from float data MAP_P_DELTA=200



Figure 15: Float 3901948. Trajectory of the Argo float plotted with historical Argo data (blue dots). The black contours indicate continent contours.



3901948 uncalibrated float data (-) and mapped salinity (o) with objective errors

Figure 16: Float 3901948. T/S diagram for the uncalibrated float data and mapped salinity. The collared circles indicate the theta levels with an objective errors. Diagram is showing every 3rd Argo profile.



3901948 potential conductivity (mmho/cm) multiplicative correction r with errors



Figure 17: Float 3901948. Potential conductivity (top) and vertically averaged salinity (bottom) with errors.



3901948 calibrated float data (-) and mapped salinity (o) with objective errors





Figure 19: Float 3901948. Salinity anomaly on Theta





Figure 20: Float 3901948. Salinities with errors on θ .



Figure 21: Float 3901948. Calibrated salinity anomaly on $\theta.$



Figure 22: Float 3901948. Salinity, salinity variance on theta and OW chosen levels.

3.3 Summary and Conclusions

4 Final Checks



Figure 23: Float 3901948. Time series of applied pressure corrections.



Figure 24: Float 3901948. Time series of applied temperature corrections.



Figure 25: Float 3901948. Time series of applied salinity corrections.