

Delayed Mode Quality Control of Argo float 3901500

Kamila Walicka

British Oceanographic Data Centre (BODC), National Oceanography Centre
Joseph Proudman Building, 6, Brownlow St, Liverpool L3 5DA

March 13, 2020

Short Summary

The pressure sensor propagate well with QC=1, err 2.4 dbar. From profiles 1 to 60 float is behaving well QC=1 error =0.01. From profiles 61 to 127 float is drifting and OWC correction has to be applied.

WMO number	DM correction
3901500	Drift detected

Table 1: Correction applied in delayed mode.

Contents

1	Introduction	3
2	Quality Check of Argo Float Data	3
2.1	Time Series of Vertical Distribution of Data	3
2.2	Comparison between Argo Float and Climatology	6
2.3	Satellite Altimeter comparison	9
3	Pressure Adjustment for APEX Floats	9
4	Correction of Salinity Data	11
4.1	Comparison between Argo floats and CTD Climatology	11
4.1.1	Configuration	11
4.1.2	Results	14
4.2	Comparison between Argo floats and Argo Climatology	22
4.2.1	Configuration	22
4.2.2	Results	26
4.3	Summary and Conclusions	34
5	References	34

1 Introduction

Delayed mode analysis was performed for float number 134522 (3901500) where salinity and temperature values were separately compared to nearby historical CTD profiles and nearby Argo profiles as a reference database. The OWC (Cabanes et al., 2016) method was run to estimate a salinity offset and a salinity drift. For more information about float 134522 (3901500) click on the following link: <http://www.ifremer.fr/argoMonitoring/float/3901500>

2 Quality Check of Argo Float Data

2.1 Time Series of Vertical Distribution of Data




Figure 1: Float 3901500. Time series of the vertical distribution of potential temperature (°C).





Figure 2: Float 3901500. Time series of the vertical distribution of salinity (PSS-78).

2.2 Comparison between Argo Float and Climatology



6

Figure 3: Float 3901500. Potential temperature ($^{\circ}$ C) plotted with pressure (dBar) and data from WMO boxes of CTD reference data (CTD for DMQC 2019V01) +/- 10 $^{\circ}$ of latitude and longitude. The black and blue cycles indicates the first and the last Argo profile, respectively. Green symbols represent other Argo profiles from this float. The thin colors lines indicate the reference data




Figure 4: Float 3901500. Salinity (PSS-78) plotted with pressure (dBar) and data from WMO boxes of CTD reference data (CTD for DMQC 2019V01) +/- 10° of latitude and longitude. The black and blue cycles indicates the first and the last Argo profile, respectively. Green symbols represent other Argo profiles from this float. The thin colors lines indicate the reference data.

Float 3901500: theta vs salinity
(symbols = float profiles,% curves = historical CTDs)




Figure 5: Float 3901500. T/S diagram plotted with and data from WMO boxes of CTD reference data (CTD for DMQC 2019V01) +/- 10° of latitude and longitude. The black and blue cycles indicates the first and the last Argo profile, respectively. Green symbols represent other Argo profiles from this float.

2.3 Satellite Altimeter comparison




Figure 6: Float 3901500. The comparison between the Sea Surface Height (SSH) from the satellite altimetry and Dynamic Height Anomaly (DHA) extracted from the Argo float temperature and salinity data (<ftp://ftp.ifremer.fr/ifremer/argo/etc/argo-ast9-item13-AltimeterComparison/figures/>).

3 Pressure Adjustment for APEX Floats

Float 3901500 is the Apex float, where the pressure sensor is not auto-corrected to zero while at the sea surface, hence the pressure data in Apex float have to be corrected during processing in delayed-mode. The procedures of adjusting sea surface pressure are described in Argo User's Manual, 2017 (<https://archimer.ifremer.fr/doc/00228/33951/32470.pdf>). The pressure sensor in Apex float 3901500 is not truncated, QC=1, error = 2.4 dbar (Figure 7).

**Raw surface pressure measured before descent (+0 dbar offset) for float
134522**
pressure correction in green





Figure 7: Float 3901500. Sea surface pressure data. The red cross indicate the raw pressure before float descent, recorded after sending data to GDAC. Blue circle indicate pressure value in the real-time. Green rotated cross shows the pressure correction applied from the previous float cycle.

4 Correction of Salinity Data

4.1 Comparison between Argo floats and CTD Climatology

4.1.1 Configuration

```
% =====
%
%     Climatology Data Input Paths
%

HISTORICAL_DIRECTORY=/users/argo/climatology
HISTORICAL_CTD_PREFIX=/historical_ctd/CTD_for_DMQC_2019V01/ctd_
HISTORICAL_BOTTLE_PREFIX=/historical_bot/WOD2001_v2/bot_
HISTORICAL_ARGO_PREFIX=/argo_profiles/ARGO_for_DMQC_2019V03/argo_

% =====
%
%     Float Input Path
%

FLOAT_SOURCE_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_source/
FLOAT_SOURCE_POSTFIX=.mat

% =====
%
%     Mapping Output Path
%

FLOAT_MAPPED_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_mapped/ctd/
FLOAT_MAPPED_PREFIX=map_
FLOAT_MAPPED_POSTFIX=.mat

% =====
%
%     Calibration Output Path
%

FLOAT_CALIB_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_calib/ctd/
FLOAT_CALIB_PREFIX=cal_
FLOAT_CALSERIES_PREFIX=calseries_
FLOAT_CALIB_POSTFIX=.mat

% =====
```

```

% Diagnostic Plots Output Path
%
FLOAT_PLOTS_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_plots/ctd/
%
% =====
%
% Constants File Path
%
CONFIG_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/constants/
CONFIG_COASTLINES=coastdat.mat
CONFIG_WMO_BOXES=wmo_boxes_ctd.mat
CONFIG_SAF=TypicalProfileAroundSAF.mat
%
% =====
%
% max number of historical casts used in objective mapping
CONFIG_MAX_CASTS=310
%
% 1=use PV constraint, 0=don't use PV constraint, in objective mapping
MAP_USE_PV=1
%
% 1=use SAF separation criteria, 0=don't use SAF separation criteria, in objective mapping
MAP_USE_SAF=1
%
% spatial decorrelation scales, in degrees
MAPSCALE_LONGITUDE_LARGE=6
MAPSCALE_LONGITUDE_SMALL=3
MAPSCALE_LATITUDE_LARGE=4
MAPSCALE_LATITUDE_SMALL=2
%
% cross-isobath scales, dimensionless, see BS(2005)
MAPSCALE_PHI_LARGE=0.1
MAPSCALE_PHI_SMALL=0.02
%
% temporal decorrelation scale, in years
MAPSCALE_AGE=10
MAPSCALE_AGE_LARGE=20
%
% exclude the top xxx dbar of the water column
MAP_P_EXCLUDE=100
%
% only use historical data that are within +/- yyy dbar from float data
MAP_P_DELTA=200

```


4.1.2 Results




Figure 8: Float 3901500. Location of the float profiles (red line with coloured numbers) and the CTD reference data selected for mapping (blue dots). The black contours indicate the bathymetry at 0, 200, 1000 and 2000 m.

3901500 uncalibrated float data (-) and mapped salinity (o) with objective errors





Figure 9: Float 3901500. Plots the original float salinity and the objectively estimated reference salinity at the 10 float theta levels that are used in calibration.

3901500 potential conductivity (mmho/cm) multiplicative correction r with errors



3901500 vertically-averaged salinity (PSS-78) additive correction




Figure 10: Float 3901500. Evolution of the suggested adjustment with time. The top panel plots the potential conductivity multiplicative adjustment. The bottom panel plots the equivalent salinity additive adjustment. The red line denotes one-to-one profile fit that uses the vertically weighted mean of each profile. The red line can be used to check for anomalous profiles relative to the optimal fit.

3901500 calibrated float data (-) and mapped salinity (o) with objective errors




Figure 11: Float 3901500. Plots of calibrated float salinity and the objectively estimated reference salinity at the 10 float theta levels that are used in calibration.




Figure 12: Float 3901500. Salinity anomaly on theta levels.





Figure 13: Float 3901500. Plots of the evolution of salinity with time along with selected theta levels with minimum salinity variance.



20

Figure 14: Float 3901500. Calibrated salinity anomaly on theta levels.







Figure 15: Float 3901500. Plots including the theta levels chosen for calibration: Top left: Salinity variance at theta levels. Top right: T/S diagram of all profiles of Argo float. Bottom left: potential temperature plotted against pressure. Bottom right: salinity plotted against pressure.

4.2 Comparison between Argo floats and Argo Climatology

4.2.1 Configuration

```
% =====
%
%     Climatology Data Input Paths
%
HISTORICAL_DIRECTORY=/users/argo/climatology
HISTORICAL_CTD_PREFIX=/historical_ctd/CTD_for_DMQC_2019V01/ctd_
HISTORICAL_BOTTLE_PREFIX=/historical_bot/bot_
HISTORICAL_ARGO_PREFIX=/argo_profiles/ARGO_for_DMQC_2019V03/argo_-
%
% =====
%
%     Float Input Path
%
FLOAT_SOURCE_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_source/
FLOAT_SOURCE_POSTFIX=.mat
%
% =====
%
%     Mapping Output Path
%
FLOAT_MAPPED_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_mapped/argo/
FLOAT_MAPPED_PREFIX=map_
FLOAT_MAPPED_POSTFIX=.mat
%
% =====
%
%     Calibration Output Path
%
FLOAT_CALIB_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_calib/argo/
FLOAT_CALIB_PREFIX=cal_
FLOAT_CALSERIES_PREFIX=calseries_
FLOAT_CALIB_POSTFIX=.mat
%
% =====
%
%     Diagnostic Plots Output Path
%
```

```

FLOAT_PLOTS_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/float_plots/argo/
% =====
% Constants File Path
%
CONFIG_DIRECTORY=/users/argo/ow/matlabow-2.0.1/data/constants/
CONFIG_COASTLINES=coastdat.mat
CONFIG_WMO_BOXES=wmo_boxes_argo.mat
CONFIG_SAF=TypicalProfileAroundSAF.mat

% =====
% Objective Mapping Parameters
%

% max number of historical casts used in objective mapping
CONFIG_MAX_CASTS=310

% 1=use PV constraint, 0=don't use PV constraint, in objective mapping
MAP_USE_PV=1

% 1=use SAF separation criteria, 0=don't use SAF separation criteria, in objective mapping
MAP_USE_SAF=1

% spatial decorrelation scales, in degrees
MAPSCALE_LONGITUDE_LARGE=6
MAPSCALE_LONGITUDE_SMALL=3
MAPSCALE_LATITUDE_LARGE=4
MAPSCALE_LATITUDE_SMALL=2

% cross-isobath scales, dimensionless, see BS(2005)
MAPSCALE_PHI_LARGE=0.1
MAPSCALE_PHI_SMALL=0.02

% temporal decorrelation scale, in years
MAPSCALE_AGE=10
MAPSCALE_AGE_LARGE=20

% exclude the top xxx dbar of the water column
MAP_P_EXCLUDE=100

% only use historical data that are within +/- yyy dbar from float data
MAP_P_DELTA=200

```


4.2.2 Results




Figure 16: Float 3901500. Location of the float profiles (red line with coloured numbers) and the CTD reference data selected for mapping (blue dots). The black contours indicate the bathymetry at 0, 200, 1000 and 2000 m.

3901500 uncalibrated float data (-) and mapped salinity (o) with objective errors





Figure 17: Float 3901500. Plots the original float salinity and the objectively estimated reference salinity at the 10 float theta levels that are used in calibration.

3901500 potential conductivity (mmho/cm) multiplicative correction r with errors



3901500 vertically-averaged salinity (PSS-78) additive correction




Figure 18: Float 3901500. Evolution of the suggested adjustment with time. The top panel plots the potential conductivity multiplicative adjustment. The bottom panel plots the equivalent salinity additive adjustment. The red line denotes one-to-one profile fit that uses the vertically weighted mean of each profile. The red line can be used to check for anomalous profiles relative to the optimal fit.

3901500 calibrated float data (-) and mapped salinity (o) with objective errors





Figure 19: Float 3901500. Plots of calibrated float salinity and the objectively estimated reference salinity at the 10 float theta levels that are used in calibration.



30

Figure 20: Float 3901500. Salinity anomaly on theta levels.




Figure 21: Float 3901500. Plots of the evolution of salinity with time along with selected theta levels with minimum salinity variance.




Figure 22: Float 3901500. Calibrated salinity anomaly on theta levels.







Figure 23: Float 3901500. SPlots including the theta levels chosen for calibration: Top left: Salinity variance at theta levels. Top right: T/S diagram of all profiles of Argo float. Bottom left: potential temperature plotted against pressure. Bottom right: salinity plotted against pressure.

4.3 Summary and Conclusions

5 References